

30. **(Reiterated)** The method of claim 28, wherein said morphogen comprises residues 330-431 of SEQ ID NO: 2.
31. **(Reiterated)** The method of claim 28, wherein said morphogen comprises residues 48-292 of SEQ ID NO: 2.
32. **(Reiterated)** The method of claim 28, wherein said morphogen comprises the amino acid sequence of SEQ ID NO: 2.
46. **(Amended)** The method of claim 28, wherein said morphogen comprises residues 293-329 of SEQ ID NO: 2.
47. **(Amended)** The method of claim 28, wherein said morphogen comprises residues 293-431 of SEQ ID NO: 2.
48. **(Reiterated)** The method of claim 28, wherein said morphogen comprises residues 30-431 of SEQ ID NO: 2.

Please add the following new claims:

52. **(New)** A method for enhancing the formation and development of dendrites and synapses in hippocampal neurons, comprising contacting said neurons with a morphogen selected from: an OP-1 polypeptide, a BMP-2 polypeptide, a BMP-5 polypeptide, a BMP-6 polypeptide, or a 60A polypeptide, wherein said morphogen has a conserved C-terminal seven-cysteine skeleton at least about 70% homologous to residues 330-431 of human OP-1 (SEQ ID NO: 2), and wherein said morphogen induces dendrite outgrowth in said hippocampal neuron.
53. **(New)** The method of claim 52, wherein said morphogen comprises residues 30-292 of SEQ ID NO: 2.
54. **(New)** The method of claim 52, wherein said morphogen comprises residues 330-431 of SEQ ID NO: 2.

55. (New) The method of claim 52, wherein said morphogen comprises residues 48-292 of SEQ ID NO: 2.
56. (New) The method of claim 52, wherein said morphogen comprises the amino acid sequence of SEQ ID NO: 2.
57. (New) The method of claim 52, wherein said morphogen comprises residues 293-329 of SEQ ID NO: 2.
58. (New) The method of claim 52, wherein said morphogen comprises residues 293-431 of SEQ ID NO: 2.
59. (New) The method of claim 52, wherein said morphogen comprises residues 30-431 of SEQ ID NO: 2.

The claims presented above incorporate changes as indicated by the marked-up versions below.

28. (Amended Twice) A method for enhancing the formation and development of dendrites and synapses in hippocampal neurons ~~cells~~, comprising contacting said neurons ~~cells~~ with a morphogen selected from: an OP-1 polypeptide, a BMP-2 polypeptide, a BMP-5 polypeptide, a BMP-6 polypeptide, ~~and~~ or a 60A polypeptide, wherein said morphogen has a conserved C-terminal seven-cysteine skeleton at least about 60% identical to residues 330-431 of human OP-1 (SEQ ID NO: 2), and wherein said morphogen induces dendrite outgrowth in said hippocampal neuron.
46. (Amended) The method of claim 28, wherein said morphogen comprises residues ~~292-330~~ 293-329 of SEQ ID NO: 2.
47. (Amended) The method of claim 28, wherein said morphogen comprises residues ~~292-431~~ 293-431 of SEQ ID NO: 2.